

PNAS

^aCenter for Computational and Evolutionary Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China; ^bDepartment of Biology, University College London, London WC1E 6BT, United Kingdom; and ^cGenome Center and Department of Evolution and Ecology, University of California, Davis, CA 95616

EVOLUTION

PNAS Early Edition | 1 of 6

There are five possible species delimitations: $\{a, b, c\}$ with one species, $\{a\}\{b, c\}$, $\{b\}\{c, a\}$, $\{c\}\{a, b\}$, each with two species, and $\{a\}\{b\}\{c\}$ with three species. The last delimitation has three different species phylogenies. The total number of possible species trees is thus seven (Fig. S1). In general, the total number of species trees for a set of individuals is much larger than the number of possible trees conditioned on a particular species delimitation (e.g., ref. 12).

If both Λ and S are treated as unknown, a full Bayesian approach would generate the joint posterior distribution of species delimitations and species trees

$$f(S, \Lambda | D) = \frac{1}{f(D)} f(D | S) f(S | \Lambda) f(\Lambda),$$

where D denotes multilocus sequence data for a sample of individuals, $f(D | S)$ is the likelihood of the data given the species phylogeny obtained by integrating over gene trees as outlined in (13), and $f(S | \Lambda)$ and $f(\Lambda)$ are prior distributions on species phylogenies and species delimitations, respectively. To infer only the species delimitations, for example, one could integrate the posterior probability with respect to the possible species trees (i.e., average over uncertainties in species trees). One advantage of this approach is that it can incorporate various sources of information regarding species delimitations. Namely, the prior $f(\Lambda)$ may be informative and based on previously observed patterns of population substructure. If prior information is lacking, $f(\Lambda)$ could be specified by assuming, for example, the Dirichlet process commonly used in Bayesian clustering (e.g., ref. 15). Similarly, $f(S | \Lambda)$ can be specified either by using prior phylogenetic information from other sources or by assigning equal probabilities to rooted trees, or labeled histories, for the species. The prior on the divergence times in the species tree specifies the amount of time that genetic isolation must have persisted before we recognize genetically isolated samples as distinct species (rather than subpopulations). This is an essential component of a species definition.

Species Delimitation Using a Guide Phylogeny. The full Bayesian approach to species delimitation outlined herein is very challenging to implement, even though all its components are computable. Here we develop a simplified Bayesian approach relying on a user-specified guide phylogeny to reduce the space of phylogenies and species delimitations that we must integrate over. The guide tree represents the phylogenetic relationships among the most subdivided possible delimitation of individuals into species that appears biologically plausible. It may be generated based on morphological characters or on geographic areas where the individuals are sampled (see below). Let S_G and Λ_G be a guide phylogeny and species delimitation, respectively. The total number of possible species delimitations, Z , depends on the specific form of the guide tree. An entirely unbalanced guide tree of s species has $Z = s$, but Z can be much greater. The following algorithm calculates Z for any guide phylogeny S :

1. Label all tips of the tree with value 1.
2. Move to the ancestor of each pair of labeled nodes and set the label value for the ancestral node to be $x \times y + 1$, where x and y are the label values of the two daughter nodes, respectively.
3. Repeat step 2 until the root node is labeled.
4. Set Z to be the value of the root node label.

For example, application of this algorithm to the guide tree of Fig. 1A gives $Z = 7$. Let S_i and Λ_i for $i \in (1, \dots, Z)$ be a species phylogeny and species delimitation, respectively, obtained by collapsing one or more nodes on the guide tree. We now construct a reversible-jump Markov chain Monte Carlo (rjMCMC) algorithm that successively splits or joins nodes on the guide




Fig. 1. Given the guide species tree (A), each species delimitation is represented by a set of flags indicating whether each of the four ancestral nodes (6, 7, 8, 9) is collapsed (0) or resolved (1). For this guide tree, there exist seven species delimitations, shown in B, where 0000 indicates all nodes are collapsed so that there is only one species, and 1111 indicates the fully resolved tree with five species. The reversible-jump algorithm allows moves between species delimitations connected in B. The probabilities of the species delimitations under the uniform Dirichlet prior with equal probabilities for each labeled history are shown in parentheses. For example, tree 1101 has prior probability 0.2 because this tree corresponds to two labeled histories, with node 9 being older or younger than node 7, respectively (node 8 is collapsed in tree 1101). The prior with equal probabilities for the rooted trees assigns probability 1/7 for each of these species delimitations.

species tree, generating the posterior probabilities for different collapsed subtrees, S_i , of the guide species tree that correspond to specific hypotheses regarding species delimitations, Λ_i . Given the guide tree, S_i is thus both a species tree and a species delimitation model. Here, we adopt the biological species concept, recognizing groups that have experienced no recent gene flow as potential species (although not requiring other evidence of reproductive isolation). Other species concepts can be accommodated in this framework, however, by modifying the priors to allow for limited hybridization, etc.

Posterior Probability of a Species Delimitation. The posterior probability of species delimitation S_i is

$$f(S_i, \Lambda_i | D, S_G, \Lambda_G) = \frac{1}{f(D)} \times \sum_{G \in \mathcal{G}} \int_{\theta} \int_{\tau_0} \int_{\tau} f(S_i, \Lambda_i | S_G, \Lambda_G) \prod_j f(D_j | G_j) \times f(G_j | \theta, S_i, \tau, \tau_0) f(\theta | S_i) \times f(\tau | S_i, \tau_0) f(\tau_0 | S_i) d\theta d\tau d\tau_0, \quad [1]$$

where D represents sequences for L loci with D_j to be the sequences at locus j , $G = \{G_j\}$, where G_j is the gene tree at locus j , τ_0 is the time of the first divergence event (at the root) on the species phylogeny, $\tau = (\tau_1, \dots, \tau_{s_i-2})$ is a vector of $s_i - 2$ nonroot node ages (in units of expected mutations per site), and s_i is the number of species in species delimitation Λ_i . Let $\theta = (\theta_1, \dots, \theta_{2s_i-2})$ be a vector of contemporary and ancestral population parameters, where $\theta_j = 4N_j\mu$, N_j is the effective population size of species j and $2s_i - 2$ is the number of branches in the species delimitation tree S_i . Note that parameter θ_j is not defined if j is a contemporary species for which one or no sequences are sampled at each locus. The likelihood $f(D_j | G_j)$ is calculated using standard methods (16) under the Jukes–Cantor mutation model (17). The prior probability density of gene trees conditional on the species tree, $f(G_j | \theta, S_i, \tau, \tau_0)$, is calculated using the equations in ref. 13.

Two priors were implemented for the species delimitation models (Fig. 1). The first assigns equal probabilities to rooted species trees. This is the default in the program and used in analysis of this paper. The second assigns equal probabilities to all labeled histories that are compatible with the guide species tree or its collapsed subtrees (18, 19). On a large unbalanced

guide tree with many potential species, this prior assigns much greater probabilities to resolved trees than to collapsed trees, and may thus be inappropriate.

Given the species tree and root age τ_0 , the rank-ordered species divergence times have the uniform Dirichlet prior

$$f(\tau|S_i, \tau_0) = (s_i - 2)! \tau_0^{-(s_i - 2)}, \quad [2]$$

where $s_i - 2$ is the number of divergence times for the nonroot nodes. The root age is assigned a gamma prior $\tau_0 \sim G(\alpha, \beta)$. The θ_j 's are independently and identically distributed according to another gamma distribution.

Reversible-Jump Markov Chain Monte Carlo. The MCMC algorithm described in ref. 9 is used with the addition of a new pair of moves that either expand or collapse a node in the guide tree. The moves are between models of different dimensions, and are implemented using rjMCMC (20).

Split. Suppose there are x previously collapsed nodes in the guide tree that may be split. A joined node is feasible for splitting if either it is the root, or its mother node is already split. Choose one of the x nodes at random for splitting. Let it be node i , and its two daughter nodes in the guide tree be j and k . The split move changes the current species-delimitation model S with parameter θ_i to a new delimitation S^* with parameters $\theta_i^*, \tau_i^*, \theta_j^*, \theta_k^*$; other parameters are shared between the two models. Note that the new species divergence time τ_i^* is constrained by the gene trees because sequences from two different species cannot coalesce until they reach their common ancestral species. The upper-bound τ_U is thus determined by scanning all gene trees to find the most recent coalescence event between two sequences of which one has ancestor j and the other has ancestor k . Our algorithm for doing this goes through all tips of the gene tree and moves toward the root, flagging each node that has j or k as ancestors. The ages of nodes that have both j and k as ancestors are used to determine τ_U . Also, τ_U should be younger than the age of the mother node on the species tree.

We tried several reversible-jump proposals, and identified two that seem to work well (algorithms 0 and 1). We describe algorithm 0 in detail and then algorithm 1 only briefly. Algorithm 0 generates three random variables, u_1 , u_2 , and u_3 , to achieve dimension matching from parameters $(\theta_i, u_1, u_2, u_3)$ in S to $(\theta_i^*, \tau_i^*, \theta_j^*, \theta_k^*)$ in S^* , as follows:

$$\begin{aligned} \theta_i^* &= \theta_i, \\ \tau_i^* &= u_1 \sim P(\tau_U), \\ \theta_j^* &= \theta_i e^{(u_2 - 0.5)}, \\ \theta_k^* &= \theta_i e^{(u_3 - 0.5)}, \end{aligned} \quad [3]$$

where u_1 is from a parabola distribution with density $f(u_1; \tau_U) = 3u_1^2/\tau_U^3$ and cumulative distribution function $F(u_1; \tau_U) = (u_1/\tau_U)^3$, $0 < u_1 < \tau_U$, and where u_2, u_3 are $U(0, 1)$ random numbers, and ϵ is a fine-tuning parameter. The move makes use of the expectation that the new τ_i should be close to the upper-bound τ_U , which reflects constraints of the gene trees (Fig. S2).

The acceptance ratio for the move is

$$\begin{aligned} R_{\text{split}} &= \frac{x}{y} \frac{\pi(S^*)g(u^*)}{\pi(S)g(u)} \times \left| \frac{\partial(\theta_i^*, \tau_i^*, \theta_j^*, \theta_k^*)}{\partial(\theta_i, u_1, u_2, u_3)} \right| \\ &= \frac{x}{y} \frac{\pi(S^*)}{\pi(S)} \frac{\tau_U^3}{3u_1^2} (\epsilon\theta_j^*)(\epsilon\theta_k^*), \end{aligned} \quad [4]$$

where x is the number of feasible nodes for splitting in S , and y is the number of feasible nodes for joining in S^* , $\pi(S^*)/\pi(S)$ is the product of the prior ratio and the likelihood ratio, and $g(u)$ and $g(u^*)$ are the probability densities for generating random variables in the source and target. The factor $(\epsilon\theta_j^*)(\epsilon\theta_k^*)$ is due to the

two new parameters (θ_j and θ_k) in S^* . However, if node j is a tip on the guide tree with at most one sequence at any locus, θ_j will not be a new parameter. The factor is replaced by $(\epsilon\theta_j^*)$ or $(\epsilon\theta_k^*)$ if θ_j only or θ_k only is created by the move, or by 1 if the move does not create any new θ parameter.

We use node IDs on gene trees to keep track of the population/species within which each coalescent event occurs, so that a node with ID i represents a coalescent that occurred in species i . With the creation of τ_i^* , we scan the gene trees to update the node IDs: if a node has ID i but its age is younger than τ_i^* , the ID is changed into j or k (for the daughter species on the guide tree).

Join. The move for merging (joining) a pair of species proceeds as follows: identify the number, x , of possible nodes on the guide tree that may be joined; a node can be joined if its two immediate descendants are either tips or joined nodes. Choose one of them with equal probability. Let this be i and its immediate descendants be j and k . Change all node IDs j and k on gene trees into i . Parameters θ_j and θ_k , if they exist, are eliminated from the model, as is parameter τ_i . Dimension matching is achieved through

$$(\theta_i, \tau_i, \theta_j, \theta_k) \rightarrow (\theta_i^*, u_1^*, u_2^*, u_3^*).$$

The acceptance ratio is

$$R_{\text{join}} = \frac{x}{y} \frac{\pi(S^*)}{\pi(S)} \frac{3u_1^{*2}}{\tau_U^3} \times \frac{1}{(\epsilon\theta_j^*)(\epsilon\theta_k^*)}, \quad [5]$$

where x is the number of feasible nodes for joining in S , and y is the number of feasible nodes for splitting in S^* ; τ_U is the upper bound for splitting node i in the target state S^* . Similarly to the split move, the factor $(\epsilon\theta_j^*)(\epsilon\theta_k^*)$ is used only if both θ_j and θ_k exist in species delimitation S .

Algorithm 1 proposes the new parameters θ_j and θ_k in the split move from a gamma distribution based on the current θ_i :

$$\begin{aligned} \theta_i^* &= \theta_i, \\ \tau_i^* &= u_1 \sim P(\tau_U), \\ \theta_j^* &= u_2 \sim G(\alpha, \alpha/(m\theta_i)), \\ \theta_k^* &= u_3 \sim G(\alpha, \alpha/(m\theta_i)), \end{aligned} \quad [6]$$

where u_1 is from the parabola distribution as described (Fig. S2), and where u_2 and u_3 are gamma variables with shape α and mean $m\theta_i$, with α and m to be fine-tuning parameters. The join move simply drops the extra parameters, as before. The acceptance ratios are

$$\begin{aligned} R_{\text{split}} &= \frac{x}{y} \frac{\pi(S^*)}{\pi(S)} \frac{\tau_U^3}{3u_1^2} \frac{1}{g(u_2; \alpha, \alpha/(m\theta_i))g(u_3; \alpha, \alpha/(m\theta_i))}, \\ R_{\text{join}} &= \frac{x}{y} \frac{\pi(S^*)}{\pi(S)} \frac{3u_1^{*2}}{\tau_U^3} \times g(u_2^*; \alpha, \alpha/(m\theta_i^*))g(u_3^*; \alpha, \alpha/(m\theta_i^*)), \end{aligned} \quad [7]$$

where $g(u; \alpha, \beta)$ is the gamma density. Similarly, the factors $g(u_2)g(u_3)$ and $g(u_2^*)g(u_3^*)$ are used only if both parameters θ_j and θ_k exist in the split tree (in which node i is resolved).

In some cases, the algorithms did not mix well in analyzing large informative datasets. In particular, an ϵ too small in algorithm 0 may result in poor mixing, because the proposed values θ_j^* and θ_k^* in the split move may be far away from the mode of the posterior. It is advisable to use a large ϵ (such as 10 or 20) and to run the same analysis at least twice using both algorithms.

Species Delimitation Without rjMCMC. For large numbers of loci and/or sequences, the rjMCMC may display mixing problems (e.g., difficulty moving between models). In such cases, a method that does not use rjMCMC may be preferable. A second method, referred to as the “ τ threshold” method for species delimitation

was therefore developed that does not require the use of rjMCMC. This approach involves integrating over only the most complex model (the fully resolved guide tree) using constant dimensional MCMC and using the posterior distribution of species divergence times to identify the species delimitations. The posterior probability P that the divergence time between a pair of putative species is below a threshold value (determined by the species definition) is interpreted as the probability that the two groups form a single species, whereas $1 - P$ is the probability that they form two distinct species. Our current implementation assigns a gamma prior on the root age τ_0 , chosen such that the prior probability of recognizing either one, or two, species at the root of the species tree is 0.5. Note that this typically favors very small τ_0 values and differs from the τ_0 prior used in the rjMCMC algorithm. The rjMCMC method may be considered similar to assigning a mixture prior on τ_0 for the fully resolved guide species tree, with a point value 0 and a gamma distribution.

Results

Statistical Performance for Simulated Data. Computer simulations were performed to examine the posterior probability associated with the correct model when the algorithm is applied to choose between the one- and two-species models. Simulations under the one-species model assumed a single-population coalescent process (21) with parameter $\theta = 4N\mu$. Simulations under the two-species model assumed independent coalescent processes in each species, both with parameter θ , until time τ in the past when the lineages enter a common ancestral population with coalescence parameter θ . Independent gene trees were simulated at each locus, and sequences 1 kb in length were simulated on each gene tree under the Jukes–Cantor (17) mutation model.

For each of six parameter combinations, 100 replicate datasets were simulated and analyzed using our program. Two sequence sample configurations were examined: (1, 1) and (5, 5), where (i, j) indicates that i sequences are sampled from one potential species, and j sequences are sampled from the other. The divergence time parameter was either $\tau = \theta$, $\tau = \theta/10$, or $\tau = 0$, the final case indicating a single species. The parameter $\theta = 0.01$ was used, corresponding to an average of 1% divergence between any pair of sequences in a single population at equilibrium. To analyze data simulated with $\tau > 0$, we assumed that the guide tree (sequence partition) was correct, whereas for data simulated with $\tau = 0$, we assumed that the sequences in each species partition were chosen at random. This corresponds to a situation in which a biologist is sampling allopatric species versus a single panmictic species. We used a gamma (1, 10) distribution (with a mean of 0.1) for both τ and θ . In both cases, this is larger than the true values used in the simulations and allowed us to examine robustness of the posterior model probabilities when the prior is misspecified to varying degrees. Each of the 600 simulated datasets was analyzed by running two independent chains (initiated with different seeds) for 10^6 iterations and checking for consistency of posterior model probabilities; the results were highly consistent between chains.

The results of the simulation study are summarized in Fig. 2A and B. First we consider the sample configuration of one sequence from each population. When the true model is the one-species model ($\tau = 0$), the posterior probability for the true (one-species model) is always greater than 90%. When the two-species model is the true model ($\tau = 0.01$ and $\tau = 0.001$), the posterior probability for the true model is typically low unless at least 10 loci are sampled. In these cases, the mean of the prior on τ is either one or two orders of magnitude larger than the true values. Nonetheless, with sufficient numbers of loci, and/or individuals, it is possible to identify the correct model. For a recent divergence between species ($\tau = 0.001$), when using a prior on τ with a much larger mean, the average posterior probability of the correct model is only about 0.70, even with a sample of 100 loci (Fig. 24). Sampling five

sequences per population dramatically improves the power of the method. In that case, if the true model is a single species ($\tau = 0$), then the posterior probability of the correct model is near 1.0 for all numbers of loci examined. If the correct model is two species with a relatively ancient divergence ($\tau = 0.01$), the posterior probability is also near 1 for all numbers of loci examined (even for a single locus), whereas for a more recent divergence ($\tau = 0.001$), the posterior probability still approaches 1 for as few as 10 loci (Fig. 2B). Overall, for the priors on τ and θ used in this study (which tend to specify values larger than the true simulation parameters), the method tends to be a species “lumper” if the power is low and the splits are generally conservative; if a species split has high posterior probability, it is very likely to be correct.

Rotifer (Genus *Rotaria*). We applied the rjMCMC method to a dataset of asexual bdelloid rotifers (22). Fontaneto et al. (22) collected *Rotaria* samples worldwide and conducted phylogenetic analysis using mitochondrial cytochrome oxidase I (COI) and nuclear 28S ribosomal genes. Using a species delimitation method based on estimated divergence times on gene trees (11), they suggested that the bdelloid rotifers formed independently evolving and distinct entities equivalent to species. The dataset consists of 77 mitochondrial COI sequences and 52 nuclear 28S sequences. Here, we analyze sequences from four traditionally recognized species: *R. tardigrada*, *R. neptunoida*, *R. sordida*, and *R. macrura*, with 28 COI and 17 28S sequences. The guide species tree is $((R. tardigrada, R. neptunoida), R. sordida), R. macrura$, shown in Fig. 3A (ref. 22, figure 3). We assign the prior $\tau_0 \sim G(2, 40)$, with mean 0.05, and $\theta \sim G(2, 200)$, with mean 0.01 (Fig. S3). Analysis of the COI data alone generates the posterior tree probability $\Pr(111) = 0.997$ and $\Pr(110) = 0.003$, where 111 represents the fully resolved tree and 110 the tree with *R. tardigrada* and *R. neptunoida* collapsed into one species. On the tree 111, the posterior means of θ s range from 0.06 to 0.18, whereas the posterior mean of the root age on the species tree is 0.077. Analysis of the 28S data alone led to $\Pr(111) = 0.750$ and $\Pr(110) = 0.248$. On the tree 111, the posterior means of θ s range from 0.01 to 0.06, and the posterior mean of the root age is 0.014. The COI sequences are much more divergent and informative than the 28S sequences. To analyze both loci, we use a Dirichlet distribution to account for variable mutation rates between loci, with $\alpha = 2$ (23, Eq. 4), obtaining $\Pr(111) = 1.000$. As bdelloid rotifers are asexual, both the COI and 28S loci are haploid.

North American Fence Lizards (*Sceloporus*). We applied the rjMCMC method to a dataset for five North American fence lizards *Sceloporus tristichus*, *S. cowlesi*, *S. consobrinus*, *S. undulates*, and *S. woodi* (24). The sample consists of 17 individuals, with 4, 3, 4, 5, and 1 individuals for the five species, respectively. There are 29 nuclear genes in the data, with the length ranging from 254 to 1,522 sites. The number of sequences at each locus range from 10 to 17 sequences. The guide tree, shown in Fig. 3B, is based on a Bayesian species tree analysis of the mitochondrial genes (mtDNA) (24). Given the guide tree, there are seven possible trees (Fig. 1B). Until recently, four of these species (*S. consobrinus*, *S. cowlesi*, *S. tristichus*, and *S. undulates*) were treated as a single polytypic species, with wide geographic distributions in the United States and central Mexico (25). The current species-level phylogeny, taxonomy, and phylogeographic assessment of the group is based on a mitochondrial DNA genealogy. Here we analyze the nuclear data to examine whether the mtDNA-based species are supported by nuclear genes.

We use the prior $\theta \sim G(2, 1000)$, with mean 0.002, and $\tau_0 \sim G(2, 1000)$, with mean 0.002 (Fig. S3). If all 29 loci are used, the fully resolved tree 1111 has posterior probability 1.00. The information concerning the species tree in this multiple-locus multiple-individual dataset seems overwhelming. The posterior means of the parameters under the model range from 0.002 to 0.005 for θ for the four extant species, and 0.001–0.003 for θ for




Fig. 2. Mean posterior probability of the correct model across 100 replicate datasets as a function of the number of unlinked loci. The sequence at each locus was 1 kb in length. In all cases $\theta = 0.01$. The divergence time $\tau = 0$ corresponds to a single species, whereas $\tau = \theta$ and $\tau = \theta/10$ correspond to two species with ancient and recent divergence times. In *A*, one sequence was sampled from each of two populations. In *B*, five sequences were sampled from each population.

the four ancestral species. The estimate of the root age τ_0 is 0.0018 with the 95% interval to be (0.0014, 0.0021). If only one locus (anonymous locus sun006) is used, the posterior probabilities are 0.54, 0.17, 0.13, 0.08, and 0.04 for trees 1111, 1101, 1110, 1001, and 1100, respectively. Even with one locus, tree 1111 reached relatively high posterior probability. This is consistent with the simulation result, which shows that the power can be high when multiple samples are taken from each species/population. The posterior for tree 1111 rises with the addition of loci, at 0.65, 0.41, 0.70, 0.97, for $L = 2-5$, for example.

Human Populations. We analyzed a dataset of human ethnic populations using the τ threshold approach in which the evidence for the populations belonging to the same species is assessed through the posterior probability that $\tau < \tau_T$, a preset threshold. We set $\tau_T = 2 \times 10^{-4}$, which means 10,000 generations of separation, based on a generation time of 20 years and a mutation rate of 10^{-9} mutations per site per year. The sequence data comprise samples from six populations (26), including three non-Africans: French Basques, Han Chinese, and Melanesians; and three Africans: Biaka from the Central African Republic, Mandenka from Senegal, and San from Namibia. The data consist of 20 autosomal loci, each of about 20 kb, with 18–32 sequences from each of the six populations (or 160 sequences in total) for each locus. We used the neighbor-joining tree constructed by Wall et al. (26) based on the F_{ST} distances between populations, shown in Fig. 3C. The same gamma prior $\theta \sim G(2, 1000)$, with mean 0.0005, is assigned to all of the 11 θ parameters. The root τ is assigned the prior $\tau_0 \sim G(1, 3500)$, with $\Pr(\tau_0 < \tau_T) = 0.50$ (Fig. S3). The prior for the four other τ s is specified by the Dirichlet distribution. The posterior probabilities $\Pr(\tau_j < \tau_T)$ are 0.98, 1.0, 1.0, 1.0, and 1.0 for nodes 7, 8, 9, 10, and

11 in the tree of Fig. 3C, respectively. Thus the analysis strongly supports the hypothesis that human individuals of all six populations are from the same species. The posterior means of the θ s range from 0.0005 for the contemporary Melanesian population to 0.012 for the ancestral population of the three African populations (node 10 of Fig. 3C).

Discussion

Impact of the Guide Tree. Here we consider a few heuristic approaches to constructing a guide tree. First, one may analyze the sequence data concatenated over loci to generate a large tree of individuals and then decide on the potential species by examining the groups defined on this tree of individuals. One may also analyze the multiple loci separately and combine the gene trees to construct a guide tree. If assignment of individuals to potential species is already accomplished, the guide tree topology may be generated using species tree methods (6, 27). Other data, such as morphological characters and geographical distributions, may also be used to construct the guide tree. Finally, the use of a few competing guide trees allows an assessment of the impact of the guide tree on the inference.

Though the use of the guide species tree has allowed us to implement the species-delimitation algorithm, it may nevertheless be a serious limitation. In our rjMCMC and τ -threshold algorithms, two individuals that are clustered into one population in the guide tree will never be separated into different species, no tree rearrangements are used to modify the guide tree, and only special cases of the guide tree (i.e., less-resolved trees generated by collapsing nodes on the guide tree) are evaluated. If the guide tree and its less-resolved special cases make up all of the species delimitations and species phylogenies that have substantial posterior probabilities, our algorithm will be a good approximation of the general algorithm outlined earlier, which considers all assignments and species trees (Λ and S). However, errors in both the assignment of individuals to populations and in the guide tree topology for the populations may cause inference errors.

For a test, the first locus in the lizard dataset was analyzed using the guide tree (((tri, con), cow), (und, woo)), which differs from the tree of Fig. 3B concerning the relationships among *Sceloporus tristichus*, *S. consobrinus*, and *S. cowlesi* (24). For easy comparison, we calculated the posterior probability that each node in the guide tree is collapsed, giving (((tri, con) 0.31, cow) 0.12, (und, woo) 0.20) 0.005, in comparison with (((tri, cow) 0.33, con) 0.12, (und, woo) 0.20) 0.004 for the guide tree of Fig. 3B. The two analyses thus gave very similar results, with probability 0.12 that the three concerned species should be lumped into one species, and probability 0.4–0.5% that all of the five species should be lumped into one species. The high similarity may be due to the fact that the two guide trees are quite similar.




Fig. 3. The guide species trees for the three empirical datasets analyzed in the text. (A) The guide tree for four bdelloid rotifer species/populations: *Rotaria tardigrada*, *R. neptunoida*, *R. sordida*, and *R. macrura*. (B) The guide tree for five lizard species/populations: *Sceloporus tristichus* (tri), *S. cowlesi* (cow), *S. consobrinus* (con), *S. undulatus* (und), and *S. woodi* (woo). (C) The guide tree for six human populations: Han Chinese (han), French Basques (bas), Melanesians (mel), Mandenka (man), Biaka (bia), and San (san).

